



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$$F(a, a, d, d) = f(a, a, d, d) f'(a, a, d, d) = 12a^2d^2 - 2a^4 - 2d^4.$$

That is, $12a^2d^2 - 2a^4 - 2d^4$ must be resolvable into two rational factors in a and d , since neither $f(a, a, d, d)$ nor $f'(a, a, d, d)$ can equal unity. It is evident however that $12a^2d^2 - 2a^4 - 2d^4$ does not possess this property.

GEOMETRY.

190. Proposed by WILLIAM HOOVER, A. M., Ph. D., Professor of Mathematics and Astronomy, Ohio University, Athens, Ohio.

Find the locus of the centers of sections of an ellipsoid by planes which are at a constant distance from the center.

Solution by the PROPOSER.

The center of the ellipsoid being the origin, and (α, β, γ) being the center of the section, its equation is found to be

$$\frac{\alpha}{a^2}x + \frac{\beta}{b^2}y + \frac{\gamma}{c^2}z - \left(\frac{\alpha^2}{a^2} + \frac{\beta^2}{b^2} + \frac{\gamma^2}{c^2} \right) = 0 \dots (1).$$

The perpendicular from the center of the ellipsoid upon it is

$$\left(\frac{\alpha^2}{a^2} + \frac{\beta^2}{b^2} + \frac{\gamma^2}{c^2} \right) \div \sqrt{\left(\frac{\alpha^2}{a^4} + \frac{\beta^2}{b^4} + \frac{\gamma^2}{c^4} \right)} = k,$$

a constant, by the problem. This gives the required locus, which, by rationalizing, is easily seen to be a surface of the fourth degree.

Excellent solutions were also received from PROFESSORS ZERR, WALKER, and SCHEFFER.

CALCULUS.

151. Proposed by J. SCHEFFER, A. M., Hagerstown, Md.

Integrate the differential equation, $xy \frac{\partial^2 z}{\partial x \partial y} = bx \frac{\partial z}{\partial x} + ay.$

Solution by WILLIAM HOOVER, A. M., Ph. D., Professor of Mathematics and Astronomy, Ohio University, Athens, Ohio.

Let $x = e^u$, $y = e^v$; then with $x \frac{d}{dx} = \theta$, $y \frac{d}{dy} = \theta'$, the given equation reduces to $\theta(\theta' - b)z = ae^v \dots (1)$, in which u and v are the independent variables.

The integral of (1) is